


ADA018391



TECHNICAL/SCIENTIFIC
REPORT



5-75

TREATMENT OF ACUTE ENTERIC FEVER WITH AMOXYCILLIN

By

Z. Farid, J.E. Sippel, A. Hassan, B. Trabolssi

U. S. NAVAL MEDICAL RESEARCH UNIT No. 3
(CAIRO, ARAB REPUBLIC OF EGYPT)
FPO NEW YORK 09527

**Best
Available
Copy**

TREATMENT OF ACUTE ENTERIC FEVER WITH AMOXYCILLIN

SIR,—With the appearance of strains of *Salmonella typhi* resistant to chloramphenicol¹⁻³ there is a continuing need for a search for other antibiotics that may be effective in the treatment of acute typhoid and paratyphoid fevers.⁴ We have found ampicillin⁵ and co-trimoxazole^{6,7} to be effective; and we now report on the use of amoxycillin in the treatment of seven patients with acute enteric fever and one patient with acute brucellosis (see table).

All the patients were very sick and toxic and had a temperature of 39°C or above before treatment. Blood and stool cultures were performed before treatment, twice weekly during treatment, and for at least 3 weeks afterwards. Three patients had positive blood cultures for *S. typhi* and three were positive for *S. paratyphi A*. One patient had a positive stool culture for *S. typhi* with a rising Widal titre. The eighth patient had a positive blood culture for *Brucella melitensis*. All patients were treated with two capsules (500 mg.) amoxycillin every 8 hours (3 times daily) and treatment continued for 7 days after return of temperature to normal. The patient with acute brucellosis was treated for 3 weeks.

The seven patients with enteric fever were cured. One of these patients, though symptom-free and afebrile, had a positive blood culture for *S. paratyphi A* 10 days after ending treatment, but he did not relapse clinically and was not given further treatment. The patient with acute brucellosis became afebrile after 6 days of treatment and was symptom-free 4 days later; however, 9 weeks after treatment he had a recurrence of fever, and a blood culture was positive for *Br. melitensis*. He was successfully treated with tetracycline and streptomycin. There were no adverse side-effects from amoxycillin, and no patient had a drug rash. These preliminary results of treating enteric fever with amoxycillin are encouraging.

CASE SUMMARIES

Case no.	Sex	Age (yr.)	Days febrile before treatment	Blood culture	Stool culture	No. of days to become afebrile	Comments
1	M	15	7	<i>S. typhi</i>	..	8	Very toxic, clinical improvement within 5 days.
2	M	22	5	<i>S. typhi</i>	<i>S. typhi</i>	7	Very toxic, clinical improvement within 5 days.
3	M	18	10	<i>S. typhi</i>	<i>S. typhi</i>	10	Very toxic, clinical improvement within 7 days.
4	F	25	7	Rising Widal titre	<i>S. typhi</i>	2	Rapid clinical improvement (2 days)
5	M	22	10	<i>S. paratyphi A</i>	..	5	Rapid clinical improvement (2 days)
6	M	21	7	<i>S. paratyphi A</i>	..	10	Clinical improvement within 7 days
7	M	21	7	<i>S. paratyphi A</i>	..	4	Symptom and afebrile, but one blood culture positive 10 days after treatment
8	M	15	4	<i>Br. melitensis</i>	..	6	Low-grade fever and blood culture positive 9 wk. after treatment

Clinical improvement = disappearance of toxicity, abdominal distention, coated tongue, and headache.

and further trials are now warranted using a higher dose similar to that used with ampicillin³ (100 mg. per kg. body-weight per day).

U.S. Naval Medical Research
Unit No. 3, Cairo, Egypt,
Abbassia Fever Hospital,
Ministry of Public Health,
Cairo, Egypt.

Z. FARID
J. E. SIPPEL
A. HASSAN
B. TRABOLSI.

1. Anderson, E. S., Smith, H. R. *Br. med. J.* 1972, iii, 329.
2. Overturf, G., Marton, K. I., Mathies, A. W. *New Engl. J. Med.* 1973, 289.
3. Gonzales-Cortes, A., Bessudo, D., Saucedo-Leyva, R., Fragoso, R., Hinojosa, M., Becerril, P. *Lancet*, 1973, ii, 605.
4. Geddes, A. M., Goodell, J. A. D. *Br. med. J.* 1972, iii, 525.
5. Robertson, R. P., Wahab, M. F. A., Raasch, F. O. *New Engl. J. Med.* 1968, 278, 171.
6. Farid, Z., Hassan, A., Wahab, M. F. A., Sanborn, W. R., Kent, D. C., Yassa, A., Hathout, S. E. *Br. med. J.* 1970, iii, 323.
7. Hassan, A., Erian, M., Said, F., Sorensen, K., Sippel, J. E. *ibid.* 1973, iii, 108.

UNCLASSIFIED
Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) U.S. Naval Medical Research Unit No. 3 PPO New York 09527		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
		2b. GROUP
3. REPORT TITLE. 6 TREATMENT OF ACUTE ENTERIC FEVER WITH AMOXYCILLIN.		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) 9 Technical/Scientific report (Letter to Editor)		
5. AUTHOR(s) (First name, middle initial, last name) 10 Z./Farid, J.E./Sippel, A./Hassan, and B./Trabolsi 12/6p.		
6. REPORT DATE 1974 11/1974	7a. TOTAL NO. OF PAGES 3	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO.	8b. ORIGINATOR'S REPORT NUMBER (S) 14 NAMRU-3-TR-5-75	
d. PROJECT NO. 17 MR041-20-11-0259	9b. OTHER REPORT NO (S) (Any other numbers that may be assigned this report) Acc.983	
10. DISTRIBUTION STATEMENT Distribution of this report is unlimited		
11. SUPPLEMENTARY NOTES Published in: LANCET, March 2, 1974 p. 350		12. SPONSORING MILITARY ACTIVITY Bureau of Medicine and Surgery Department of the Navy Washington, D.C. 20372
13. ABSTRACT ➔ With the appearance of strains of <u>Salmonella typhi</u> resistant to Chloramphenicol there is a continuing need for search for other antibiotics that may be effective in the treatment of acute typhoid and paratyphoid fevers. Ampicillin and co-trimoxazole were found to be effective in the treatment of typhoid and paratyphoid fevers; and we now report on the use of amoxycillin in the treatment of seven patients with acute enteric fever and one patient with acute brucellosis. is reported. ↗		

UNCLASSIFIED
Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Enteric Fever Typhoid Paratyphoid Ampicillin Co-trimoxazole Amoxycillin Therapy						